

Recombinant Human CCL22/MDC Protein (His Tag)

Catalog Number:PKSH031817



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	A-152E5.1;ABCD-1;DC/B-CK;MDC;SCYA22;STCP-1
Species	Human
Expression Host	E.coli
Sequence	Gly 25-Gln 93
Accession	O00626
Calculated Molecular Weight	9.7 kDa
Observed molecular weight	10 kDa
Tag	N-His

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 30% Acetonitrile, 0.1% TFA Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Chemokine (C-C motif) ligand 22(ABCD-1 / CCL22) is a kind of CC chemokine which is a family of secreted proteins involved in immunoregulatory and inflammatory processes. The cytokine displays chemotactic activity for monocytes, dendritic cells, natural killer cells and for chronically activated T lymphocytes. It also displays a mild activity for primary activated T lymphocytes and has no chemoattractant activity for neutrophils, eosinophils and resting T lymphocytes. This ABCD-1 / CCL22 chemokine binds to chemokine receptor CCR4. This chemokine may play a role in the trafficking of activated / effector T lymphocytes to inflammatory sites and other aspects of activated T-lymphocyte physiology. ABCD-1 / CCL22 is highly expressed in macrophage and in monocyte-derived dendritic cells, and thymus, and in Langerhans' cell histiocytosis and atopic dermatitis but not in dermatopathic lymphadenopathy. This chemokine is also

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human CCL22/MDC Protein (His Tag)

Catalog Number:PKSH031817



found in lymph node, appendix, activated monocytes, resting and activated macrophages. This protein is lower expressed in lung and spleen and very weakly expressed in small intestine.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017