



A Reliable Research Partner in Life Science and Medicine

# Elab Fluor® 488 Anti-Human CD39 Antibody[A1]

Catalog Number: E-AB-F1165L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Reactivity Human Mouse Host

Mouse IgG1, κ Isotype

Clone No. A1

Isotype Control Elab Fluor® 488 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792L]

Conjugation Elab Fluor®488

**Conjugation Information** Elab Fluor® 488 is designed to be excited by the Blue laser (488 nm) and detected using

an optical filter centered near 520 nm (e.g., a 525/40 nm bandpass filter).

Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer. Storage Buffer

#### **Applications** Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount **FCM** 

> of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

# **Preparation & Storage**

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

**Shipping** Ice bag

## **Antigen Information**

**Alternate Names** ATPDase;CD 39;NTPDase1;SPG64

**Uniprot ID** P49961 Gene ID 953

**Background** Human CD39 is an integral membrane protein with two transmembrane domains. It

> exists as a homotetramer. Expression of CD39 is found on activated lymphocytes, a subset of T cells and B cells, and dendritic cells with weak staining on monocytes and granulocytes. CD39 and CD73 have been found on regulatory T cells, specifically the effector/memory like T cells. CD39 can hydrolyze both nucleoside triphosphates and diphosphates. CD39 is the dominant ecto nucleotidase of vascular and placental trophoblastic tissues and appears to modulate the functional expression of type 2 purinergic (P2) G protein coupled receptors (GPCRs). CD39 has intrinsic ecto-ATPase activity. Expression of CD39 is induced on T cells and increased on B cells as a late

activation antigen. Product Details

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Email:techsupport@elabscience.com