

## Recombinant Phospho-Erk 1, 2 (Thr202, Tyr204) Monoclonal Antibody

catalog number: **AN300713L**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Reactivity</b>	Human;Mouse;Rat
<b>Immunogen</b>	A synthetic peptide corresponding to residues around (Thr202, Tyr204) of Human Phospho-Erk 1, 2
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG,k
<b>Clone</b>	B652
<b>Purification</b>	Protein A
<b>Buffer</b>	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

### Applications Recommended Dilution

<b>WB</b>	1:2000-1:10000
<b>IF</b>	1:200-1:1000
<b>ELISA</b>	1:5000-1:20000

### Data



Western Blot with Recombinant Phospho-Erk 1, 2 (Thr202, Tyr204) Monoclonal Antibody at dilution of 1:1000 dilution.

Lane A: Hela-2 cell lysate.

**Observed-MW:44 kDa,42 kDa**

**Calculated-MW:44 kDa,42 kDa**

### Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	Ice bag

### Background

The protein encoded by this gene is a member of the MAP kinase family. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act in a signaling cascade that regulates various cellular processes such as proliferation, differentiation, and cell cycle progression in response to a variety of extracellular signals. This kinase is activated by upstream kinases, resulting in its translocation to the nucleus where it phosphorylates nuclear targets. Alternatively spliced transcript variants encoding different protein isoforms have been described.

### For Research Use Only