

Biotin Anti-Mouse CD16/32 Antibody[2.4G2]

Catalog Number: E-AB-F0997B

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Mouse
Host	Rat
Isotype	Rat IgG2b, κ
Clone No.	2.4G2
Isotype Control	Biotin Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09843B]
Conjugation	Biotin
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is $\leq 1.0 \mu\text{g}$ per 10^6 cells in 100 μL volume or 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
------------	---

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	CD16a/b;CD32;CD32A/B;FCG2A;FCGR2A/BFCGR3;FCGR3A/B;Fc fragment of IgG low affinity IIIa/b receptor;Fc fragment of IgG low affinity IIIb receptor;Fc fragment of IgG low affinity IIa/b receptor;Fc gamma RIla/bFc gamma receptor III A/B;FcGR
Uniprot ID	P08508;P08101
Gene ID	14130;14131
Background	CD16 is low affinity IgG Fc receptor III (FcR III) and CD32 is FcR II. CD16/CD32 are expressed on B cells, monocytes/macrophages, NK cells, granulocytes, mast cells, and dendritic cells. The Fc receptors bind antibody-antigen immune complexes and mediate adaptive immune responses.

For Research Use Only