



A Reliable Research Partner in Life Science and Medicine

Biotin Anti-Mouse CD16/32 Antibody[2.4G2]

Catalog Number: E-AB-F0997B

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

Isotype Rat IgG2b, κ **Clone No.** 2.4G2

Isotype Control Biotin Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09843B]

Conjugation Biotin

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. For flow

cytometric staining, the suggested use of this reagent is \leq 1.0 μ g per 10⁶ cells in 100 μ L volume or 100 μ L of whole blood. It is recommended that the reagent be titrated for

optimal performance for each application.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD16a/b;CD32;CD32A/B;FCG2A;FCGR2A/BFCGR3;FCGR3A/B;Fc fragment of IgG low

affinity Illa/b receptor;Fc fragment of IgG low affinity Illb receptor;Fc fragment of IgG low

affinity IIa/b receptor;Fc gamma RIIa/bFc gamma receptor III A/B;FcGR

 Uniprot ID
 P08508;P08101

 Gene ID
 14130;14131

Background CD16 is low affinity IgG Fc receptor III (FcR III) and CD32 is FcR II. CD16/CD32 are

expressed on B cells, monocytes/macrophages, NK cells, granulocytes, mast cells, and dendritic cells. The Fc receptors bind antibody-antigen immune complexes and

mediate adaptive immune responses.