

CCL4/MIP-1 β /ACT-2, Human, Recombinant**Cat. No. : GPCK296****产品信息**

物种	Human
表达宿主	E.coli
序列信息	APMGSDPPTACCFSYTARKLPNHFVVDYYETSSLCSQPAVVFQTKRGKQVCADPSES WVQEY VYDLELN.
检索号	P13236.1
标签	tag-free
分子量	7.75 kDa
有效期	12 months
生物活性	Measure by its ability to chemoattract BaF3 cells transfected with human CCR5. The ED50 for this effect is < 10 ng/mL.

产品特性

内毒素 (EU/ μ g)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored at -5~-20°C or -80°C for 3-6 months.
运输	Ambient temperature or ice pack.
制剂	The protein was lyophilized from a 0.2 μ m filtered solution containing 50 mM Tris and 150 mM NaCl, pH 8.5.
复溶	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μ g/mL. Do Not Vortex! Vigorous shaking may impair the biological activity of the protein.

背景介绍

CCL4 act as an important pro-inflammatory cytokine during acute and chronic inflammatory responses. CCL4 also play a role as an attractant NK cells, dendritic cells, monocytes, and lymphocytes to sites of injury and inflammation. As CCL4 conducts signaling through G-protein-coupled receptor CCR5, which is a major co-receptor for M-tropic HIV strains. Thus, binding of CCL4 to CCR5 inhibits HIV entry and reduces the cell surface expression of CCR5. CCL4 has been identified as one of the major anti-HIV factors produced by CD8+ T cells.

