

# Recombinant Human CST9L/Testatin Protein (Fc Tag)

Catalog Number:PKSH030679

 **DIA-AN®**  
by Elabscience

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	bA218C14.1;CTES7B;PRO3543;UNQ1835
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Met 1-His 147
<b>Accession</b>	Q9H4G1
<b>Calculated Molecular Weight</b>	41.3 kDa
<b>Observed molecular weight</b>	48 kDa
<b>Tag</b>	C-hFc

## Properties

<b>Purity</b>	> 92 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 92 % as determined by reducing SDS-PAGE.

## Background

Testatin is a member of the Cystatin family. Cystatins comprise genes that all show expression patterns that are strikingly restricted to reproductive tissue. Cystatins are a family of cysteine protease inhibitors with homology to chicken cystatin. There are typically about 115 amino acids in this family. They are largely acidic, contain four conserved cysteine residues known to form two disulfide bonds, may be glycosylated and/or phosphorylated, with similarity to fetuins, kininogens, stefins, histidine-rich glycoproteins and cystatin-related proteins. Testatin shows homology to family 2 cystatins, a group of broadly expressed small secretory proteins that are inhibitors of cysteine proteases in vitro but whose in vivo functions are unclear. It is expressed in germ cells and somatic cells in reproductive tissues. Testatin is considered a strong candidate for involvement in early testis development. Testatin expression is maintained in the adult Sertoli cell, and it can also be

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human CST9L/Testatin Protein (Fc Tag)

Catalog Number:PKSH030679



found in a small population of germ cells.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)