

PerCP/Cyanine 5.5 Anti-Human CD268 Antibody[6G10]

Catalog Number: AN00655J

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	6G10
Isotype Control	PerCP/Cyanine5.5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792J]
Conjugation	PerCP/Cyanine5.5
Conjugation Information	PerCP/Cyanine 5.5 is designed to be excited by the blue laser (488 nm) and detected using an optical filter centered near 675 nm (e.g., a 690/50 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications

Recommended usage

FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.
-----	---

Data



Staining of normal human peripheral blood cells with FITC Anti-Human CD19 Antibody and PerCP/Cyanine 5.5 Anti-Human CD268 Antibody[6G10] (left) or PerCP/Cyanine 5.5 Mouse IgG1, κ Isotype Control (right). Cells in the lymphocytes gate were used for analysis.

Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 24 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

Antigen Information

Alternate Names	TNFRSF13C;BAFF-R;BAFFR;BR3;BAFF Receptor;CD268
Uniprot ID	Q96RJ3
Gene ID	115650

For Research Use Only

Background

B cell-activating factor receptor (BAFF-R) is a 19 kD type III membrane protein. It belongs to TNFR superfamily, also known as TNFRSF member 13C (TNFRSF13C), BAFF receptor 3 (BR3), or CD268. BAFF-R is expressed on mature B cells, B cell lymphoma, and T cell subset. BAFF-R is the major receptor for BAFF/BLys (or TALL-1, THANK) which binds to TACI and BCMA as well. The interaction of BAFF with BAFF-R promotes NF- κ B activation and plays predominant roles in B-cell maturation and survival as well as costimulates T cell activation and proliferation. TRAF3 is a BAFF-R intracellularly associated protein, which negatively regulates BAFF-R-mediated NF- κ B activation.