

**IL-4/BSF-1 (C-6His), Mouse, Recombinant****Cat. No. : GPCK034****产品信息**

物种	Mouse
表达宿主	Human Cells
序列信息	His21-Ser140
检索号	P07750
标签	C-6His
分子量	14.6 kDa
有效期	12 months
生物活性	Measured in a cell proliferation assay using M-NFS-60 mouse lymphoblast cells. The ED50 for this effect is 104 pg/mL.

**产品特性**

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 μm filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4.
复溶	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**背景介绍**

Interleukin-4 (IL-4) is a pleiotropic Cytokine that regulates diverse T and B cell responses including cell proliferation, survival and gene expression. IL-4 is produced by mast cells, T cells, and bone marrow stromal cells. IL-4 regulates the differentiation of naive CD4<sup>+</sup> T cells into helper Th2 cells, characterized by their Cytokine-secretion profile that includes secretion of IL-4, IL-5, IL-6, IL-10, and IL-13, which favor a humoral immune response. Another dominant function of IL-4 is the regulation of immunoglobulin class switching to the IgG1 and IgE isotypes. Excessive IL-4 production by Th2 cells has been associated with elevated IgE production and allergic response.

