## Recombinant Human Interleukin-2/IL-2 Protein (His Tag)

Catalog Number: PKSH033425




Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Mol_Mass	16.4 kDa
Accession	P60568
<b>Bio-activity</b>	Measure by its ability to induce proliferation in CTLL-2 cells. The $\mathrm{ED}_{50}$ for this effect
	is $<3$ ng/mL. The specific activity of recombinant human IL-2 is $>5$ x $10^6$ IU/mg.
Properties	

	is <3 ng/mL. The specific activity of recombinant numan IL-2 is >5 x 10° 10/mg.
Properties	
Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	$< 0.01 \; EU \; per \; \mu g \; of the protein as determined by the LAL method.$
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS,pH 8.0.
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

## Data



> 98 % as determined by reducing SDS-PAGE.

## Background

Interleukin-2(IL-2) is an interleukin; a type of cytokine signaling molecule in the immune system; belongs to the IL-2 family. It is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. IL-2/IL-2R signaling is required for T-cell proliferation and other fundamental functions that are essential for the immune response. IL-2 stimulates growth and differentiation of B-cells; NK cells; lymphokine-activated killer cells; monocytes; macrophages and oligodendrocytes.

## For Research Use Only