

TTC4 Polyclonal Antibody

catalog number: E-AB-18699

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


| | |
|---------------------|--|
| Reactivity | Human;Mouse |
| Immunogen | Fusion protein of human TTC4 |
| Host | Rabbit |
| Isotype | IgG |
| Purification | Antigen affinity purification |
| Conjugation | Unconjugated |
| buffer | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |

Applications

Recommended Dilution

| | |
|------------|------------|
| IHC | 1:40-1:200 |
|------------|------------|

Data



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using TTC4 Polyclonal Antibody at dilution of 1:45($\times 200$)

Preparation & Storage

| | |
|-----------------|--|
| Storage | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. |
| Shipping | The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended. |

Background

This gene encodes a protein that contains tetratricopeptide (TPR) repeats, which often mediate protein-protein interactions and chaperone activity. The encoded protein interacts with heat shock proteins 70 and 90. Alternative splicing results in multiple transcript variants. Naturally-occurring readthrough transcription occurs from upstream gene MROH (maestro heat-like repeat family member 7) to this gene.

For Research Use Only