Recombinant Mouse Coagulation Factor X/F10 Protein (His Tag)

Catalog Number: PKSM040986



Note: Centrifuge before opening to ensure complete recovery of vial contents.

| - | esc | | | | | . • | | |
|-----|-----|----|---|-----|---|-----|---|---|
| - 1 | Δ | Q. | | 141 | n | ш | n | m |
| ய | U | 0 | U | ш | v | UΙ | v | ш |

Species Mouse

 Mol_Mass
 34.6&18.4 kDa

 Accession
 O88947

Bio-activity Not validated for activity

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin $< 1.0 \text{ EU} \text{ per } \mu\text{g} \text{ of the protein as determined by the LAL method.}$

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of 20mM MES, 150mM NaCl, 1mM CaCl₂

, pH 7.5.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Mouse coagulation factor X/F10 a member of the peptidase S1 family. The mature F10 is composed mostly of two EGF-like domains, one Gla gamma-carboxy-glutamate domain and one peptidase S1 domain. Factor Xa is a vitamin K-dependent plasma protease that converts prothrombin to thrombin in the presence of factor Va, calcium and phospholipid during blood clotting. The two chains of F10 are formed from a single-chain precursor by the excision of two Arg residues. A single-chain precursor is initially synthesized in the liver. The light and heavy chains are linked together by disulfide bonds. The light chain contains a Gla and two EGF-like domains. The heavy chain corresponds to the serine protease domain. It can form a heterodimer with SERPINA5.

For Research Use Only