

A Reliable Research Partner in Life Science and Medicine

# Elab Fluor® Violet 450 Anti-Human CD19 Antibody[CB19]

Catalog Number: E-AB-F1004Q

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Reactivity Human Host Mouse

**Isotype** Mouse IgG1, κ

Clone No. CB19

Isotype Control Elab Fluor<sup>®</sup> Violet 450 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-

F09792Q1

**Conjugation** Elab Fluor<sup>®</sup> Violet 450

**Conjugation Information** Elab Fluor<sup>®</sup> Violet 450 is designed to be excited by the violet laser (405 nm) and

detected using an optical filter centered near 450 nm (e.g., a 450/45 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.


#### Applications Recommended usage

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

#### **Data**



Human pheripheral blood cells are stained with Elab Fluor® Violet 450 Anti-Human CD19 Antibody (filled gray histogram). Unstained pheripheral blood cells (blank black histogram) are used as control.

## **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

### **Antigen Information**

Alternate Names B-lymphocyte antigen CD19;CD19;Cd19;Differentiation antigen CD19

Web: www.elabscience.cn

 Uniprot ID
 P15391

 Gene ID
 930

## For Research Use Only



## Elabscience Biotechnology Co., Ltd.

A Reliable Research Partner in Life Science and Medicine

## **Background**

CD19 is a 95 kD type I transmembrane glycoprotein also known as B4. It is a member of the immunoglobulin superfamily expressed on B-cells (from pro-B to blastoid B cell s, absent on plasma cells) and follicular dendritic cells. CD19 is involved in B cell development, activation, and differentiation. CD19 forms a complex with CD21 (CR2) and CD81 (TAPA-1), and functions as a BCR co-receptor.