



A Reliable Research Partner in Life Science and Medicine

Purified Anti-Mouse H-2 Antibody[M1/42]

catalog number: E-AB-F1216A

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

 Isotype
 Rat IgG2a, κ

 Clone
 M1/42

Conjugation Unconjugated


Buffer Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer.

Dialyze to completely remove the stabilizer prior to labeling.

Applications Recommended Dilution

FCM $2 \mu g/mL(0.5 \times 10^6 - 1 \times 10^6 \text{ cells})$

Data



C57/BL6 Mouse splenocytes were stained with 0.2 µg Purified Anti-Mouse H-2 Antibody[M1/42] (Right) and 0.2 µg Rat IgG2a, κ Isotype Control (Left), followed by FITC-conjugated Goat Anti-Rat IgG Secondary Antibody.

Preparation & Storage

Storage Storage, avoid freeze /

thaw cycles.

Shipping Ice bag

Background

The M1/42 antibody reacts with the H-2 MHC class I alloantigens expressed on nucleated cells from mice of the a, b, d, j, k, s, and u haplotypes (Stallcup, KC et al, 1981). MHC class I is involved in antigen presentation to T cells expressing CD3/TCR and CD8 proteins.

Email: techsupport@elabscience.cn