# Recombinant Cavia porcellus CTLA-4/CD152 Protein (His Tag)

Catalog Number: PKSQ050093



Note: Centrifuge before opening to ensure complete recovery of vial contents.

		crip				
	00	CI	РΤ	n	tπ	Λn
JU.	$\mathbf{c}$	v.		w	ш	UП

SpeciesCavia porcellusMol\_Mass16-35 kDaAccessionH0VUB1

**Bio-activity** Immobilized Mouse B7-1-Fc at 5 μg/ml (100 μl/well) can bind Cavia porcellus CTLA-

4-His. The ED<sub>50</sub> of Recombinant Cavia porcellus CTLA-4-His is 4.29 ng/ml.

## **Properties**

**Purity** > 90 % as determined by reducing SDS-PAGE.

**Endotoxin**  $< 1.0 \text{ EU per } \mu\text{g of the protein as determined by the LAL method.}$ 

Storage Storage Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.


**Shipping** This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel

packs. Upon receipt, store it immediately at < - 20°C.

**Formulation** Supplied as a 0.2 μm filtered solution of PBS, pH7.4.

**Reconstitution** Not Applicable

### Data



> 90 % as determined by reducing SDS-PAGE.

## Background

Cytotoxic Tlymphocyte 4(CTLA-4,CD152), is a type I transmembrane T cell inhibitory molecule that is a member of the Ig superfamily.CD28 and CTLA-4, together with their ligands, B7-1 and B7-2, constitute one of the dominant costimulatory pathways that regulate T and B cell responses. CD28 and CTLA-4 are structurally homologous molecules that are members of the immunoglobulin (Ig) gene superfamily. CTLA4 transmits an inhibitory signal to T cells, whereas CD28 transmits a stimulatory signal. Intracellular CTLA4 is also found in regulatory T Cells and may play an important role in their functions. Tcell activation through the Tcell receptor and CD28 leads to increased expression of CTLA4.

#### For Research Use Only