## 普诺赛® Procell system

## IGF-II, Mouse, Recombinant

Cat. No.: GPCK338

## 产品信息

**物种** Mouse

表达宿主 E.coli

序列信息 AYGPGETLCGGELVDTLQFVCSDRGFYFSRPSSRANRRSRGIVEECCFRSCDLALLETYCATPA

cellsystem

KSE with polyhistidine tag at the N-terminus.

**检索号** P09535.1

标签 His-tag at the N-terminus

**分子量** 8.20 kDa **有效期** 12 months

生物活性 Measure by its ability to induce MCF-7 cells proliferation. The ED50 for this effect is < 6

ng/mL. The specific activity of recombinant mouse IGF-II is > 1.5 × 10^5 IU/mg.

产品特性

**内毒素 (EU/μg)** < 0.1

保存 Lyophilized protein should be stored at -5~-20°C for 1 year. Upon reconstitution, store at 2-

8°C for up to 1 week. Further dilute in a buffer containing a carrier protein or stabilizer (e.g. 0.1% BSA, 10% FBS, 5% HSA or 5% trehalose solution), protein aliquots should be stored

at -5~-20°C or -80°C for 3-6 months.

运输 Ambient temperature or ice pack.

制剂 The protein was lyophilized from a 0.2 µm filtered solution containing 1 × PBS, pH 8.0.

复溶 It is recommended to reconstitute the lyophilized protein in sterile water to a

concentration not less than 100 µg/mL. Do Not Vortex! Vigorous shaking may impair the

biological activity of the protein.

## 背景介绍

Insulin-like growth factor 2 (IGF-2) is one of three protein hormones that share structural similarity to insulin. The major role of IGF-2 is as a growth promoting hormone during gestation. It is believed to be a major fetal growth factor in contrast to Insulin-like growth factor 1, which is a major growth factor in adults.

网站: <u>www.procell.com.cn</u> 电话: 400-999-2100

邮箱: techsupport@procell.com.cn

地址:湖北省武汉市高新大道858号生物医药产业园三期C4栋



