

A Reliable Research Partner in Life Science and Medicine

Recombinant VPAC2 Monoclonal Antibody

catalog number: AN301688L

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Rat; Mouse

Immunogen Recombinant human VPAC2 fragment

HostRabbitIsotypeIgG, κCloneA391

Purification Protein A purified


Buffer PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution

IHC 1:200-1:500

IF 1:50


Data



Immunohistochemistry of paraffin-embedded Human tonsil using VPAC2 Monoclonal Antibody at dilution of 1:500. Immunohistochemistry of paraffin-embedded Mouse spleen using VPAC2 Monoclonal Antibody at dilution of 1:500.



Immunohistochemistry of paraffin-embedded Rat spleen using VPAC2 Monoclonal Antibody at dilution of 1:500.



Immunofluorescent analysis of (4% Paraformaldehyde) fixed Jurkat cells using anti-VPAC2 Monoclonal Antibody at dilution of 1:50.

Preparation & Storage

Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping Ice bag

Background

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com
 Rev. V1.0

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Human VIPR2 (Vasoactive Intestinal Polypeptide Receptor 2), also known as VPAC2, is a 60-70 kDa seven transmembrane glycoprotein receptor, which belongs to the G protein-coupled receptor 2 family of proteins. In its first extracellular region hormone receptor domain (aa 49-114) contains a mature chain, which has three N-linked glycosylation sites and a "Cilia targeting signal", which is composed of and Arg -Composition of Asp-Tyr-Arg sequence. Human VIPR2 has relatively high homology with mouse and rat VIPR2. VIPR2 is mainly expressed in skeletal muscle, CD4+ T cells, smooth muscle cells, dendritic cells, liver precursor cells, single cells, synthetic cells and selective neurons. VIPR2 couples with CAMP-mediated signal transduction pathways, and binds VIP and pituitary adenase activating polypeptide (PACAP).

For Research Use Only