
Recombinant AK2/Adenylate kinase 2 Monoclonal Antibody


catalog number: AN300119P

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Human
Immunogen	Recombinant Human AK2 protein
Host	Rabbit
Isotype	IgG
Clone	9D8
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS
Applications	Recommended Dilution
WB	1:500-1:2000
IHC-P	1:100-1:500

Data





Immunohistochemistry of paraffin-embedded human liver using AK2 / Adenylate kinase 2 Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded human kidney using AK2 / Adenylate kinase 2 Monoclonal Antibody at dilution of 1:200.

Western Blot with AK2 / Adenylate kinase 2 Monoclonal Antibody at dilution of 1:500. Lane A: Mouse kidney tissue lysate, Lane B: HCT116 Whole Cell Lysate, Lysates/proteins

> at 30 μg per lane. **Observed-MW:26 kDa**

Calculated-MW:26 kDa

Preparation & Storage		
Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.	
Shipping	Ice bag	
Background		
For Research Use Only		
Toll-free: 1-888-852-8623 Web: <u>w w w .elabscience.com</u>	Tel: 1-832-243-6086 Email:techsupport@elabscience.com	Fax: 1-832-243-6017

Elabscience®

Elabscience Bionovation Inc. A Reliable Research Partner in Life Science and Medicine

Adenylate kinases are involved in regulating the adenine nucleotide composition within a cell by catalyzing the reversible transfer of phosphate groups among adenine nucleotides. Three isozymes of adenylate kinase, namely 1, 2, and 3, have been identified in vertebrates; this gene encodes isozyme 2. Expression of these isozymes is tissue-specific and developmentally regulated. Isozyme 2 is localized in the mitochondrial intermembrane space and may play a role in apoptosis. Mutations in this gene are the cause of reticular dysgenesis. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 1 and 1.