

A Reliable Research Partner in Life Science and Medicine

Recombinant AK2/Adenylate kinase 2 Monoclonal Antibody

catalog number: AN300119P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human AK2 protein

Host Rabbit
Isotype IgG
Clone A1233
Purification Protein A

Buffer 0.2 µm filtered solution in PBS


Applications Recommended Dilution

WB 1:500-1:2000 **IHC-P** 1:100-1:500


Data



Immunohistochemistry of paraffin-embedded human liver using AK2 / Adenylate kinase 2 Monoclonal Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded human kidney using AK2 / Adenylate kinase 2 Monoclonal Antibody at dilution of 1:200.



Western Blot with AK2 / Adenylate kinase 2 Monoclonal Antibody at dilution of 1:500. Lane A: Mouse kidney tissue lysate, Lane B: HCT116 Whole Cell Lysate, Lysates/proteins at 30 µg per lane.

Observed-MW:26 kDa Calculated-MW:26 kDa

Preparation & Storage

Storage This antibody can be stored at 2°C-8°C for one month without detectable loss of

activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

Shipping Ice bag

Background

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: w w w .elabscience.com
 Email: techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Adenylate kinases are involved in regulating the adenine nucleotide composition within a cell by catalyzing the reversible transfer of phosphate groups among adenine nucleotides. Three isozymes of adenylate kinase, namely 1, 2, and 3, have been identified in vertebrates; this gene encodes isozyme 2. Expression of these isozymes is tissue-specific and developmentally regulated. Isozyme 2 is localized in the mitochondrial intermembrane space and may play a role in apoptosis. Mutations in this gene are the cause of reticular dysgenesis. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 1 and 1.

For Research Use Only