A Reliable Research Partner in Life Science and Medicine

# Recombinant Mouse CD300a/LMIR1 Protein (Fc Tag)

Catalog Number: PKSM041101

Note: Centrifuge before opening to ensure complete recovery of vial contents.

$\mathbf{r}$		crip					
	00	0	РΤ	n	ŤΤ		m
v	$\mathbf{c}$	v.		w	w	w	ш

**Species** Mouse

Source HEK293 Cells-derived Mouse CD300a/LMIR1 protein Leu28-Arg183, with an C-

terminal Fc

Calculated MW 44.3 kDa
Observed MW 58-75 kDa
Accession Q6SJQ0

**Bio-activity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

LMIR1, also termed CD300a, is a type I transmembrane glycoprotein with a single IgV-like extracellular domain and an extended membrane proximal region that links the immunoglobulin (Ig) and transmembrane domains and belongs to the immunoglobulin superfamily. The intracellular domain of LMIR1 contains several immunoreceptor tyrosine-based inhibition motifs (ITIMs). When cross-linked, it will be tyrosine phosphorylated and capable of recruiting tyrosine phosphatases (SHP-1, SHP-2) and inositol polyphosphate 5-phosphatase, SHIP. LMIR1 will regulate mast cell-mediated inflammatory responses. LMIR1 is broadly expressed on myeloid and lymphoid cells, and its expression is differentially regulated depending on the cell type.

### For Research Use Only