

## Recombinant Human RPS19 Protein

**Catalog Number:** PKSH032021

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human RPS19 protein Pro2-His 145
<b>Calculated MW</b>	16.1 kDa
<b>Observed MW</b>	16 kDa
<b>Accession</b>	P39019
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, 1mM EDTA, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

40S Ribosomal Protein S19 (RPS19) is a ribosomal protein that belongs to the ribosomal protein S19e family. RPS19 is located in the nucleoli, and higher level expression is seen in colon carcinoma tissue than normal colon tissue. It required for pre-rRNA processing and maturation of 40S ribosomal subunits. RPS19 plays a role in many biological processes, such as endocrine pancreas development, erythrocyte differentiation, mRNA metabolic process. Defects in RPS19 are the cause of Diamond-Blackfan anemia type 1 (DBA 1), which is a form of Diamond-Blackfan anemia, a congenital non-regenerative hypoplastic anemia that usually presents early in infancy. Diamond-Blackfan anemia is characterized by a moderate to severe macrocytic anemia, erythroblastopenia, and an increased risk of malignancy.

### For Research Use Only