Elabscience®

Human PGRN Antibody Pair Set

Catalog No.	E-KAB-0517	Applications	ELISA
Synonyms	proepithelin;PC cell-derived growth f	actor;	

Kit components & Storage

Title	Specifications	Storage
Human PGRN Capture Antibody	1 vial, 100 µ g	Store at -20° C for one year. Avoid
		freeze/thaw cycles.
Human PGRN Detection Antibody	1 vial, 50 μL	Store at -20°C for one year. Avoid
(Biotin)		freeze/thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0517)	
		Human PGRN Capture Antibody	Human PGRN Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Human PGRN protien	Recombinant Human PGRN protien
Information	Swissprot	P28799	
Product details	Reactivity	Human	Human
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50%	PBS with 0.04% Proclin 300; 1%
		glycerol; pH 7.5	protective protein; 50% glycerol; pH
			7.5
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Human PGRN in ELISAs.	

For Research Use Only

Elabscience®

Applications

Human PGRN Sandwich ELISA Assay

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4 μg/mL	Human PGRN Capture	
Capture		Antibody	10
			Alistae
ELISA	1:1000-1:10000	Human PGRN Detection	Objeal Density
Detection		Antibody (Biotin)	· · · · · · · · · · · · · · · · · · ·
			0.01 10 100 1000 10000
			Human PGRN Concentration (pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Secreted protein that acts as a key regulator of lysosomal function and as a growth factor involved in inflammation , wound healing and cell proliferation.Secreted protein that acts as a key regulator of lysosomal function and as a growth factor involved in inflammation , wound healing and cell proliferation.Facilitates also the acidification of lysosomes , causing degradation of mature CTSD by CTSB.In addition , functions as a wound-related growth factor that acts directly on dermal fibroblasts and endothelial cells to promote division , migration and the formation of capillary-like tubule structures .

For Research Use Only