

## Human PCX Antibody Pair Set

<b>Catalog No.</b>	E-KAB-0460	<b>Applications</b>	ELISA
<b>Synonyms</b>	GCTM-2 antigen;Gp2;PC;PCLP;PCLP-1;PCLP1;PODXL;Podocalyxin-like protein 1		

### Kit components & Storage

Title	Specifications	Storage
Human PCX Capture Antibody	1 vial, 100 μg	Store at -20°C for one year. Avoid freeze/thaw cycles.
Human PCX Detection Antibody (Biotin)	1 vial, 50 μL	Store at -20°C for one year. Avoid freeze/thaw cycles.


**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Product Information

Items		Characteristic (E-KAB-0460)	
		Human PCX Capture Antibody	Human PCX Detection Antibody (Biotin)
Immunogen Information	Immunogen	Recombinant Human PCX protien	Recombinant Human PCX protien
	Swissprot	O00592	
Product details	Reactivity	Human	Human
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5 mg/mL	/
	Buffer	PBS with 0.04% Proclin 300; 50% glycerol; pH 7.5	PBS with 0.04% Proclin 300; 1% protective protein; 50% glycerol; pH 7.5
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Human PCX in ELISAs.	

## Applications

Human PCX Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images
ELISA Capture	0.5-4 µg/mL	Human PCX Capture Antibody	
ELISA Detection	1:1000-1:10000	Human PCX Detection Antibody (Biotin)	

**Note:** This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

## Background

Involved in the regulation of both adhesion and cell morphology and cancer progression. Functions as an anti-adhesive molecule that maintains an open filtration pathway between neighboring foot processes in the podocyte by charge repulsion. Acts as a pro-adhesive molecule, enhancing the adherence of cells to immobilized ligands, increasing the rate of migration and cell-cell contacts in an integrin-dependent manner. Induces the formation of apical actin-dependent microvilli. Involved in the formation of a preapical plasma membrane subdomain to set up initial epithelial polarization and the apical lumen formation during renal tubulogenesis. Plays a role in cancer development and aggressiveness by inducing cell migration and invasion through its interaction with the actin-binding protein EZR. Affects EZR-dependent signaling events, leading to increased activities of the MAPK and PI3K pathways in cancer cells.