Elabscience®

GTF3A Polyclonal Antibody

catalog number: E-AB-18063

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse;Rat
Immunogen	Synthetic peptide of human GTF3A
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Conjugation	Unconjugated
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
IHC	1:50-1:300
Data	
Immunohistochemistry tissue using GTF3A I	of paraffin-embedded Human tonsil Polyclonal Antibody at dilution of 1:75(×200)
Immunohistochemistry	Polyclonal Antibody at dilution of 1:75(×200)
Immunohistochemistry tissue using GTF3A I	Polyclonal Antibody at dilution of 1:75(×200)
Immunohistochemistry tissue using GTF3A I Preparation & Storage	Polyclonal Antibody at dilution of 1:75(×200)

Background

The product of this gene is a zinc finger protein with nine Cis[2]-His[2] zinc finger domains. It functions as an RNA polymerase III transcription factor to induce transcription of the 5S rRNA genes. The protein binds to a 50 bp internal promoter in the 5S genes called the internal control region (ICR), and nucleates formation of a stable preinitiation complex. This complex recruits the TFIIIC and TFIIIB transcription factors and RNA polymerase III to form the complete transcription complex. The protein is thought to be translated using a non-AUG translation initiation site in mammals based on sequence analysis, protein homology, and the size of the purified protein.