

A Reliable Research Partner in Life Science and Medicine

Elab Fluor® 700 Anti-Mouse CD127/IL-7RA Antibody[A7R34]

Catalog Number: E-AB-F1023M1

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Host Rat

Isotype Rat IgG2a, κ **Clone No.** A7R34

Isotype Control Elab Fluor[®] 700 Rat IgG2a, κ Isotype Control[2A3] [Product E-AB-F09832M1]

Conjugation Elab Fluor® 700

Conjugation Information Elab Fluor[®] 700 is designed to be excited by the Red laser (627-640 nm) and detected

using an optical filter centered near 719 nm (e.g., a 725/40 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Staining of C57BL/6 murine splenocytes with FITC Anti-

Mouse CD3 Antibody[17A2] and Elab Fluor[®] 700 Anti-Mouse CD127/IL-7RA Antibody[A7R34] (left) or Elab Fluor[®] 700 Rat IgG2a, κ Isotype Control (right). Total viable cells were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping lce bag

Antigen Information

Alternate Names CD127;IL-7 receptor subunit alpha;IL-7R subunit alpha;IL-7R-alpha;IL-7RA;II7r;

Interleukin-7 receptor subunit alpha

Uniprot ID P16872

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Gene ID **Background** 16197

CD127 is a 60-90 kD type I transmembrane glycoprotein also known as IL-7 receptor α chain or IL-7Rα. It forms a heterodimer with the common γ chain (γc or CD132) which is shared with the receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and IL-21. CD127 is expressed on immature B cells through early pre-B stage, thymocytes (except CD4/ CD8 double positive thymocytes), peripheral T cells, and bone marrow stromal cells. CD127 has been reported to be an useful marker for identifying memory and effector T cells. The ligation of IL-7 with its receptor is important for stimulation of mature and immature T cells as well as immature B cells proliferation and development.

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web:www.elabscience.com

Email:techsupport@elabscience.com