Recombinant Human NPRC/NPR3 (C-Fc)

Catalog Number: PKSH034022

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Human HEK293 Cells-derived Human NPRC;NPR3 protein Thr24-Glu481, with an C-terminal Fc 77.5 kDa 90-100 kDa 917342 Not validated for activity > 95 % as determined by reducing SDS-PAGE.	
Fc 77.5 kDa 90-100 kDa P17342 Not validated for activity > 95 % as determined by reducing SDS-PAGE.	
77.5 kDa 90-100 kDa P17342 Not validated for activity > 95 % as determined by reducing SDS-PAGE.	
90-100 kDa P17342 Not validated for activity > 95 % as determined by reducing SDS-PAGE.	
P17342 Not validated for activity > 95 % as determined by reducing SDS-PAGE.	
Not validated for activity > 95 % as determined by reducing SDS-PAGE.	
> 95 % as determined by reducing SDS-PAGE.	
< 1.0 EU per µg of the protein as determined by the LAL method.	
Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of	
reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	
This product is provided as lyophilized powder which is shipped with ice packs.	
Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.	
Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants	
before lyophilization.	
Please refer to the specific buffer information in the printed manual.	
Please refer to the printed manual for detailed information.	

Data

kDa	мк	R
120		
90		-
60	_	
40		
30	-	-

> 95 % as determined by reducing SDS-PAGE.

Background

Atrial Natriuretic Peptide Receptor-3 (NPR3), also known as NPRC or ANPR-C, is one of the three natriuretic peptide receptors, is a type I transmembrane glycoprotein. The natriuretic system is key to the maintenance of vascular tone and cardiovascular homeostasis. Receptor for the natriuretic peptide hormones, binding with similar affinities atrial natriuretic peptide NPPA/ANP, brain natriuretic peptide NPPB/BNP, and C-type natriuretic peptide NPPC/CNP. May function as a clearance receptor for NPPA, NPPB and NPPC, regulating their local concentrations and effects. Osteocrin was found to be a specific ligand to NPR3. NPR3 is necessary for Osteocrin to regulate femoral, tibial, and metatarsal bone elongation.

For Research Use Only Toll-free: 1-888-852-8623 Web:w w w.elabscience.com