Recombinant Bovine TGFBR1 protein (His Tag)

Catalog Number: PDEB100010

Note: Centrifuge before opening to ensure complete recovery of vial contents.

	Description			
Calculated MW20.6 kDaObserved MW20 kDaAccessionO46680Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Species	Bovine		
Observed MW20 kDaAccessionO46680Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Source	E.coli-derived Bovine TGFBR1 protein Ile143-Leu330, with an N-terminal His		
AccessionO46680Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Calculated MW	20.6 kDa		
Bio-activityNot validated for activityPropertiesPurity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Observed MW	20 kDa		
Properties Purity > 95% as determined by reducing SDS-PAGE. Endotoxin < 10 EU/mg of the protein as determined by the LAL method Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. Shipping This product is provided as lyophilized powder which is shipped with ice packs. Formulation Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol. Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of	Accession	O46680		
Purity> 95% as determined by reducing SDS-PAGE.Endotoxin< 10 EU/mg of the protein as determined by the LAL method	Bio-activity	Not validated for activity		
Endotoxin< 10 EU/mg of the protein as determined by the LAL methodStorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.ReconstitutionIt is recommended that sterile water be added to the vial to prepare a stock solution of	Properties			
StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Purity	> 95% as determined by reducing SDS-PAGE.		
°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Endotoxin	< 10 EU/mg of the protein as determined by the LAL method		
ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.ReconstitutionIt is recommended that sterile water be added to the vial to prepare a stock solution of	Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80		
ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.ReconstitutionIt is recommended that sterile water be added to the vial to prepare a stock solution of		°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of		
FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.ReconstitutionIt is recommended that sterile water be added to the vial to prepare a stock solution of		reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.		
ReconstitutionMannitol.It is recommended that sterile water be added to the vial to prepare a stock solution of	Shipping	This product is provided as lyophilized powder which is shipped with ice packs.		
Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of	Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with 5% Trehalose and 5%		
		Mannitol.		
0.5 ms/mL. Concentration is measured by LWV Ke	Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of		
0.5 mg/mL. Concentration is measured by $0.7 vis$.		0.5 mg/mL. Concentration is measured by UV-Vis.		

Data

KDa	М	R
135	-	
100	_	
75	-	
65	-	
45		
35	-	
25	-	
		-
	-	-
15	-	

> 95 % as determined by reducing SDS-PAGE.

Background

On ligand binding, forms a receptor complex consisting of two type II and two type I transmembrane serine/threonine kinases. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators. Receptor for TGF-beta.