

## Elab Fluor® Violet 540 Anti-Human CD4 Antibody[RPA-T4]

Catalog Number: E-AB-F1109T3

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Reactivity	Human
Host	Mouse
Isotype	Mouse IgG1, κ
Clone No.	RPA-T4
Isotype Control	Elab Fluor® Violet 540 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792T3]
Conjugation	Elab Fluor® Violet 540
Conjugation Information	Elab Fluor® Violet 540 is designed to be excited by the violet laser (405 nm) and detected using an optical filter centered near 548 nm (e.g., a 572/28 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.


### Applications

### Recommended usage

#### FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. **The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood).** Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### Data



Staining of normal human peripheral blood cells with APC

Anti-Human CD3 Antibody[OKT-3] and Elab Fluor® Violet 540 Anti-Human CD4 Antibody[RPA-T4](left) or Elab Fluor® Violet 540 Mouse IgG1, κ Isotype Control(right). Cells in the lymphocytes gate were used for analysis.

### Preparation & Storage

Storage	Keep as concentrated solution. This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.
Shipping	Ice bag

### Antigen Information

Alternate Names	CD4;T-cell surface antigen T4/Leu-3;T-cell surface glycoprotein CD4
Uniprot ID	P01730

### For Research Use Only

**Gene ID**

920

**Background**

CD4, also known as T4/Leu-3, is a 55 kD single-chain type I transmembrane glycoprotein and member of the immunoglobulin superfamily. It is expressed on most thymocytes, helper T cells, type II NKT cells, and monocytes/macrophages. CD4 is part of the TCR/CD3 complex, binds to  $\beta 2$  domain from the MHC class II molecule, and participates in TCR signal transduction. CD4 is the receptor of IL-16 and is a coreceptor for the human immunodeficiency virus (HIV) and human herpes virus 7 (HHV-7).