

## Recombinant Human EIF1AX Protein (His Tag)

Catalog Number: PKSH032372

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	E.coli-derived Human EIF1AX protein Met 1-Ile144, with an N-terminal His
<b>Calculated MW</b>	18.6 kDa
<b>Observed MW</b>	22 kDa
<b>Accession</b>	P47813
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
	Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Eukaryotic Translation Initiation Factor 1A, X-Chromosomal (EIF1AX) is an essential eukaryotic translation initiation factor that belongs to the eIF-1A family. EIF1AX is required for the binding of the 43S complex (a 40S subunit, eIF2/GTP/ Met-tRNA<sub>i</sub> and eIF3) to the 5' end of capped RNA and has been shown to interact with IPO13. EIF1AX contains one S1-like domain and seems to be required for maximal rate of protein biosynthesis. Enhances ribosome dissociation into subunits and stabilizes the binding of the initiator Met-tRNA(I) to 40 S ribosomal subunits.

### For Research Use Only