

Purified Anti-Human TCRV γ 9 Antibody[B3]

Catalog Number: GF003570P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human
Immunogen	Recombinant Human TCRV γ 9 protein
Host	Mouse
Isotype	Mouse IgG1, κ
Clone	B3
Purification	>98%, Protein A/G purified
Conjugation	Unconjugated
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications

Recommended Dilution

FCM	2 μ g/mL (0.5×10^6 - 1×10^6 cells)
------------	---

Data



Human peripheral blood lymphocytes were stained with 0.2 μ g Purified Anti-Human TCRV γ 9 Antibody[B3] (Right) and 0.2 μ g Mouse IgG1, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Mouse IgG Secondary Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

The V γ 9 TCR is a variant of the TCR γ chain expressed on a subset of $\gamma\delta$ T cells. V γ 9V δ 2 T lymphocytes, a major $\gamma\delta$ T cell subset in humans, recognize phosphoantigens, certain tumor cells, and cells treated with aminobisphosphonates. This cell population displays cytolytic activity against various tumor cells. The $\gamma\delta$ TCR is a heterodimeric TCR complex composed of covalently bound γ and δ chains involved in antigen recognition and the non-covalently associated monomeric proteins CD3 δ , γ , ϵ , and ζ chains. The B3 antibody reacts specifically with human TCR V γ 9 as designated by the Lefranc/Foster nomenclature system. Human TCR V γ 9 is also known as TCR V γ 2 under the Strauss, Quertermous nomenclature system. Several TCR γ and δ chain nomenclature systems exist. In order to consolidate the various nomenclature systems, we have created charts for corresponding names across some of the most popular naming methods.