Purified Anti-Mouse CD27 Antibody[LG.3A10]

catalog number: **AN003220P**


Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Reactivity	Mouse
Immunogen	Recombinant Mouse CD27 protein
Host	Armenian Hamster
Isotype	Armenian Hamster IgG1, κ
Clone	LG.3A10
Purification	>98%, Protein A/G purified
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications	Recommended Dilution
FCM	≤ 0.2 μg per million cells in 100 μL volume

Data



C57/BL6 Mouse splenocytes cell were stained with 0.2 μg Purified Anti-Mouse CD27 Antibody[LG.3A10](Right) and 0.2 μg Armenian Hamster IgG1, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Armenian Hamster IgG Secondary Antibody, then anti-Mouse B220 FITC-conjugated Monoclonal Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

CD27 is also known as S152 and T14. A member of the tumor necrosis factor receptor (TNFR) superfamily, it is a 45 kD protein expressed on peripheral T cells, memory B cells, NK cells, and thymocyte subset. Through its ligand, CD70, CD27 plays a key role in T cell and B cell interactions. Additionally, ligation of CD27 on naïve T cells may be important in their maturation to effector cells.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com