A Reliable Research Partner in Life Science and Medicine

Recombinant Rat CD155/PVR/NECL5 Protein (His Tag)

Catalog Number: PKSR030409

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Species Rat

Source HEK293 Cells-derived Rat CD155/PVR/NECL5 protein Met 1-Gly 352, with an C-

terminal His

 Calculated MW
 35.8 kDa

 Observed MW
 55-65 kDa

 Accession
 NP_058772.2

Bio-activity Immobilized rat CD155 at 20 μg/ml (100 μl/well) can bind mouse CD226 with a linear

ranger of 6. 4-160 ng/ml.

Properties

Purity > 97 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

CD155, commonly known as PVR (poliovirus receptor) and Necl-5 (nectin-like molecule-5), is a type I transmembrane single-span glycoprotein, and belongs to the nectins and nectin-like (Necl) subfamily. CD155 was originally identified based on its ability to mediate the cell attachment and entry of poliovirus (PV), an etiologic agent of the central nervous system disease poliomyelitis. The normal cellular function is in the establishment of intercellular adherens junctions between epithelial cells. CD155 may assist in an efficient humoral immune response generated within the intestinal immune system. It has been demonstrated that CD155 can be recognized and bond by DNAM-1 and CD96 which promote the adhension, migration and NK-cell killing, and thus efficiently prime cell-mediated tumor-specific immunity.

Fax: 1-832-243-6017