

Recombinant Human p38 delta/MAPK13 Protein (GST Tag)

Catalog Number: PKSH031448

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	Baculovirus-Insect Cells-derived Human p38 delta/MAPK13 protein Met 1-Leu 365, with an N-terminal GST
Calculated MW	68.4 kDa
Observed MW	60 kDa
Accession	O15264
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20mM Tris, 500mM NaCl, 2mM GSH, 10% glycerol, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Reconstitution	Please refer to the specific buffer information in the printed manual.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

The p38 family of mitogen-activated protein kinases (MAPK) includes p38 alpha (SAPK2a, CSBP), p38 beta (SAPK2b), p38 delta (SAPK4), and p38 gamma (SAPK3/ERK6). p38 alpha and p38 beta are widely expressed p38 isoforms that are involved in regulation of cell proliferation, differentiation, development, and response to stress. p38 delta, also known as MAPK13, is a regulator of differentiation-dependent gene expression in keratinocytes, and been as a regulator of surface epithelia differentiation and apoptosis. p38 delta protein is upregulated in Cholangiocarcinoma (CC) relative to hepatocellularcarcinoma (HCC) and to normal biliary tract tissues. p38 delta is important for motility and invasion of CC cells, suggesting that p38 delta may play an important role in CC metastasis. p38 delta is expressed in the epidermis, suggesting a role for p38 delta in regulating differentiation. p38 delta is the major p38 isoform driving suprabasal involucrin gene expression and that p38 delta directly regulates ERK1/2 activity via formation of a p38 delta-ERK1/2 complex. Recent emerging evidence suggests that the p38 stress MAPK pathway may function as a tumor suppressor through regulating Ras-dependent and -independent proliferation, transformation, invasion and cell death by isoform-specific mechanisms. p38 delta has important role in promoting cell proliferation and tumor development in epidermis and may have therapeutic implication for skin cancer.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017