

Recombinant Human CEACAM8/CD66b/CD67 Protein(His Tag)

Catalog Number: GPEH0940

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	E.coli-derived Human CEACAM8 protein Gln35-Asp320, with an N-terminal His
Calculated MW	31.4 kDa
Observed MW	36 kDa
Accession	P31997
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human CEACAM8/CD66b/CD67 proteins, 2µg/lane of Recombinant Human CEACAM8/CD66b/CD67 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 36 KD.

Background

CEACAM8, also known as CD66b or NCA-95, is a single chain, GPI-anchored, highly glycosylated protein belonging to the carcinoembryonic antigen family. There are four members in this family: CD66a, CD66b, CD66c, and CD66d. Members of CEACAM family are widely expressed especially on human neutrophils, and, depending on the tissue, capable of regulating diverse functions including tumor promotion, tumor suppression, angiogenesis, and neutrophil activation. Ectopic CD66 expression is commonly detected in B-cell lineage acute lymphoblastic leukemia (ALL). CEACAM8(CD66b) is also an activation marker for human granulocytes. However, its biological functions are largely unknown in eosinophils. It has been reported that CD66b is highly expressed on the surface of human peripheral blood eosinophils isolated from healthy individuals. Engagement of CD66b by mAb or a natural ligand, galectin-3, activated a Src kinase family molecule, hemopoietic cell kinase (Hck), and induced cellular adhesion, superoxide production, and degranulation of eosinophils. CD66b molecules were localized in lipid rafts, and disruption of lipid rafts or removal of the GPI anchor inhibited the adhesion and activation of eosinophils. CD66b molecules are involved in regulating adhesion and activation of eosinophils, possibly through their localization in lipid rafts and interaction with other cell surface molecules, such as CD11b. Binding of exogenous or endogenous carbohydrate ligands(s) to CD66b may be important in the release of proinflammatory mediators by human eosinophils.