



A Reliable Research Partner in Life Science and Medicine

# **Human CHEM Antibody Pair Set**

Catalog No. E-KAB-0017 Applications ELISA

**Synonyms** RARRES2, TIG2, Retinoic acid receptor responder protein 2, Tazarotene-induced gene 2 protein

## **Kit components & Storage**

Title	Specifications	Storage
Human CHEM Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Human CHEM Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

#### **Product Information**

Items		Characteristic (E-KAB-0017)	
		Human CHEM Capture Antibody	Human CHEM Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Human CHEM protein	Recombinant Human CHEM protein
Information	Swissprot	Q99969	
Product details	Reactivity	Human	Human
	Host	Mouse	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Human CHEM in ELISAs.	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>





A Reliable Research Partner in Life Science and Medicine

## **Applications**

Human CHEM Sandwich ELISA Assav:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Human CHEM Capture Antibody	
Capture			Alig
ELISA	1:1000-1:10000	Human CHEM Detection Antibody	Optical Density
Detection		(Biotin)	0.01 0.01 1 10 100  Human CHEM concentration(ng/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

### **Background**

This gene encodes a secreted chemotactic protein that initiates chemotaxis via the ChemR23 G protein-coupled seven-transmembrane domain ligand. Expression of this gene is upregulated by the synthetic retinoid tazarotene and occurs in a wide variety of tissues. The active protein has several roles, including that as an adipokine and as an antimicrobial protein with activity against bacteria and fungi.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: <a href="mailto:www.elabscience.com">www.elabscience.com</a> Email: <a href="mailto:techsupport@elabscience.com">techsupport@elabscience.com</a>