



A Reliable Research Partner in Life Science and Medicine

Purified Anti-Human CD122 Antibody[TU27]

catalog number: E-AB-F11510P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Recombinant Human CD122 protein

Host Mouse

Isotype Mouse IgG1, κ

Clone TU27

Purification >98%, Protein A/G purified


Buffer Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer.

Dialyze to completely remove the stabilizer prior to labeling.

Applications Recommended Dilution

FCM 2 μ g/mL(0.5×10⁶-1×10⁶ cells)

Data



Human peripheral blood lymphocytes cell were stained with 0.2 μ g Purified Anti-Human CD122 Antibody[TU27] (Right) and 0.2 μ g Mouse IgG1, κ Isotype Control (Left), followed by

Elab Fluor[®] 647-conjugated Goat Anti-Mouse IgG Secondary Antibody, then anti-Human CD56 PE-conjugated Monoclonal Antibody.

Preparation & Storage

Storage Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /

thaw cycles.

Shipping Ice bag

Background

Functional IL-2 receptors can exist in two affinity states on cell surfaces, the high affinity complex consisting of heterotrimers of the alpha, beta, and gamma chains and the intermediate affinity complex comprising heterodimers of the beta and gamma chains. Individual beta chains and alpha chains exhibit low affinity IL-2 binding, and the gamma chain alone does not bind IL-2. In addition to their involvement in IL-2 mediated signal transduction, both the beta chain and gamma chain have been shown to be required for IL-15 mediated signaling. IL-2 R beta is a member of the cytokine receptor superfamily.

For Research Use Only