

## HGF R/c-Met/HGFR (C-Fc), Human, Recombinant

Cat. No. : GPCK029

### 产品信息

物种	Human
表达宿主	Human Cells
序列信息	Glu25-Thr932
检索号	P08581
标签	C-Fc
分子量	128.4 kDa
有效期	12 months

### 产品特性

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
复溶	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

### 背景介绍

Hepatocyte Growth Factor Receptor (HGF R) is a glycosylated Receptor tyrosine kinase that plays a central role in epithelial morphogenesis and cancer development. HGF R is synthesized as a single chain precursor which undergoes cotranslational proteolytic cleavage. Mature HGF R is a disulfide-linked dimer composed of a 50 kDa extracellular α chain and a 145 kDa transmembrane β chain. Proteolysis and alternate splicing generate additional forms of human HGF R which either lack of the kinase domain, consist of secreted extracellular domains, or are deficient in proteolytic separation of the α and β chains. The sema domain, which is formed by both α and β chains of HGF R, mediates both Ligand binding and Receptor dimerization. HGF stimulation induces HGF R downregulation via internalization and proteasomedependent degradation. Paracrine induction of epithelial cell scattering and branching tubulogenesis results from the stimulation of HGF R on undifferentiated epithelium by HGF released from neighboring mesenchymal cells.

网站: [www.procell.com.cn](http://www.procell.com.cn)

电话: 400-999-2100

邮箱: [techsupport@procell.com.cn](mailto:techsupport@procell.com.cn)

地址: 湖北省武汉市高新大道858号生物医药产业园三期C4栋

