A Reliable Research Partner in Life Science and Medicine

MRPS18B Polyclonal Antibody

catalog number: E-AB-52720

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human

Immunogen Fusion protein of human MRPS18B

Host Rabbit
Isotype IgG


Purification Antigen affinity purification

Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Applications Recommended Dilution

WB 1:500-1:2000 **IHC** 1:40-1:200

Data



Western blot analysis of 231 and Jurkat cell lysates using MRPS18B Polyclonal Antibody at dilution of 1:250

nmunohistochemistry of paraffin-embedded I

Immunohistochemistry of paraffin-embedded Human tonsil tissue using MRPS18B Polyclonal Antibody at dilution of 1:35(×200)

Observed-MW:Refer to figures

Calculated-MW:29 kDa

Preparation & Storage

Storage Storage Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein that belongs to the ribosomal protein S18P family. The encoded protein is one of three that has significant sequence similarity to bacterial S18 proteins. The primary sequences of the three human mitochondrial S18 proteins are no more closely related to each other than they are to the prokaryotic S18 proteins. Pseudogenes corresponding to this gene are found on chromosomes 1q and 2q.

For Research Use Only

Rev. V1.7