



A Reliable Research Partner in Life Science and Medicine

Mouse TNF-a Antibody Pair Set

Catalog No. E-KAB-0094 Applications ELISA

Synonyms DIF, TNF-alpha, TNFA, TNFSF2,TNF,TNFa

Kit components & Storage

Title	Specifications	Storage
Mouse TNF-α Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Mouse TNF-α Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0094)	
		Mouse TNF-α Capture Antibody	Mouse TNF-α Detection Antibody
			(Biotin)
Immunogen	Immunogen	Recombinant Mouse TNF-α protein	Recombinant Mouse TNF-α protein
Information	Swissprot	P06804	
Product details	Reactivity	Mouse	Mouse
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Mouse TNF-α in ELISAs.	

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com





A Reliable Research Partner in Life Science and Medicine

Applications

Mouse TNF-α Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Mouse TNF-α Capture Antibody	
Capture			Aisi
ELISA Detection	1:1000-1:10000	Mouse TNF-α Detection Antibody (Biotin)	0.01 0 0.01 1000 10000 Mouse TNF-α concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

This gene encodes a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. Members of this family are classified based on primary sequence, function, and structure. This protein is synthesized as a type-II transmembrane protein and is reported to be cleaved into products that exert distinct biological functions. It plays an important role in the innate immune response as well as regulating homeostasis but is also implicated in diseases of chronic inflammation. In mouse deficiency of this gene is associated with defects in response to bacterial infection, with defects in forming organized follicular dendritic cell networks and germinal centers, and with a lack of primary B cell follicles. Alternative splicing results in multiple transcript variants.