## AF/LE Purified Anti-Human/Mouse CD44 Antibody[IM7]

catalog number: E-AB-F11000



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human;Mouse
Immunogen	Recombinant Human; Mouse CD44 protein
Host	Rat
Isotype	Rat IgG2b, κ
Clone	IM7
Purification	>98%, Protein A/G purified
Conjugation	None (AF/LE)
buffer	Sterile PBS, pH 7.2. $< 1.0$ EU per mg of the antibody as determined by the LAL method

Applications	Recommended Dilution
FCM	$2 \ \mu g/mL(1 \times 10^5 - 5 \times 10^5 \text{ cells})$
Data	
Canit 20 30 40 50	

C57/BL6 Mouse splenocytes were stained with 0.2  $\mu$ g AF/LE Purified Anti-Human/Mouse CD44 Antibody[IM7] (Right) and 0.2  $\mu$ g Rat IgG2b,  $\kappa$  Isotype Control (Left), followed by Alexa Fluor® 647-conjugated Goat Anti-Rat IgG Secondary

Antibody.

Preparation & Storage	
Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze /
	thaw cycles. This preparation contains no preservatives, thus it should be handled
	under aseptic conditions.
Shipping	Ice bag
Deelyguound	

Background

CD44 is a 80-95 kD glycoprotein also known as Hermes, Pgp1, H-CAM, or HUTCH. It is expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As B and T cells become activated or progress to the memory stage, CD44 expression increases from low or mid levels to high levels. Thus, CD44 has been reported to be a valuable marker for memory cell subsets. High CD44 expression on Treg cells has been associated with potent suppressive function via high production of IL-10. CD44 is an adhesion molecule involved in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of inflammation, and leukocyte aggregation.

## For Research Use Only