Recombinant PSGL-1/CD162/SELPLG Monoclonal Antibody

catalog number: AN300443P


Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Reactivity	Human
Immunogen	Recombinant Human PSGL-1/CD162/SELPLG Protein
Host	Rabbit
Isotype	IgG
Clone	B369
Purification	Protein A
Buffer	0.2 μm filtered solution in PBS

Applications	Recommended Dilution
WB	1:500-1:1000
IP	0.2-1 μL/mg of lysate


Data



Immunoprecipitation analysis using 1 μL anti-SELPLG Monoclonal Antibody and 15 μl of 50 % Protein G agarose. Western blot was performed from the immunoprecipitate using SELPLG Monoclonal Antibody at a dilution of 1:500. Lane A:0.5 mg Jurkat Whole Cell Lysate

Observed-MW:120 kDa

Calculated-MW:43 kDa



Western Blot with SELPLG Monoclonal Antibody at dilution of 1:500 dilution. Lane A: Jurkat Whole Cell Lysate, Lysates/proteins at 30 μg per lane.

Observed-MW:120 kDa

Calculated-MW:43 kDa

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

P-selectin glycoprotein ligand-1 (PSGL-1), also known as SELPLG or CD162, is the high affinity ounter-receptor for P-selectin on expressed on activated endothelial cells and platelets. PSGL-1 is a mucin-type glycoprotein, expressed on leukocytes and platelets as a homodimer of two disulfide-linked subunits of ~12 kD. As cell adhesion molecules, multiple studies have shown that PSGL-1/ P-selectin interaction is required for the normal recruitment of leukocytes during inflammatory reactions, and also participates in hemostatic responses. PSGL-1 protein requires two distinct posttranslational modifications for the Ca2+-dependent recognition by the lectin domain of P-selectin, that is tyrosine sulfation and specific O-linked glycosylation (sialic acid and fucose). PSGL-1 can also bind to other two members of the selectin family, E-selectin (endothelial) and L-selectin (leukocyte), but binds best to P-selectin.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com