# Recombinant Human PIGR/SC protein (His Tag)

Catalog Number: PDEH100808



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Mol Mass	59.6 kDa

 Mol\_Mass
 59.6 kDa

 Accession
 P01833

**Bio-activity** Not validated for activity

#### **Properties**

Description

**Purity** > 95% as determined by reducing SDS-PAGE.

**Endotoxin** < 10 EU/mg of the protein as determined by the LAL method

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

**Reconstitution** It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

## Data



> 95 % as determined by reducing SDS-PAGE.

## Background

Polymeric immunoglobulin receptor, also known as &nbsp,PIGR, is a member of the immunoglobulin superfamily &nbsp, and a &nbsp,Fc receptor. The ectodomain of this receptor consists of five units with homology to the variable units of immunoglobulins and a transmembrane region, which also has some homology to certain immunoglobulin variable regions. PIGR is expressed on several glandular epithelia including those of liver and breast. The deduced amino-acid sequence has a length of 764 residues and shows an overall similarity of 56% and 64% with the rabbit and rat counterpart. PIGR mediates transcellular transport of polymeric immunoglobulin molecules, and thus facilitates the secretion of &nbsp,IgA &nbsp,and &nbsp,IgM. During this process, a cleavage occurs that separates the extracellular (known as the secretory component) from the transmembrane segment of PIGR.

#### For Research Use Only