

## Recombinant Rhesus macaque B7-H4/VTCN1 Protein (His Tag)

Catalog Number: PKSQ050083

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Rhesus macaque
<b>Source</b>	HEK293 Cells-derived Rhesus macaque B7-H4/VTCN1 protein Phe29-Ala258, with an C-terminal His
<b>Calculated MW</b>	26.4 kDa
<b>Observed MW</b>	40-60 kDa
<b>Accession</b>	F7B770
<b>Bio-activity</b>	Loaded Recombinant Rhesus Macaque B7-H4 on HIS1K Biosensor, can bind Human B7-H4 mAb with an affinity constant of 0.578 pM as determined in BLI assay.

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
	Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

B7 Homolog 4 (B7-H4) is glycosylated member of the B7 family of immune costimulatory proteins. It is widely expressed, including in kidney, liver, lung, pancreas, placenta, prostate, spleen, testis and thymus. B7-H4 negatively regulates T-cell mediated immune response by inhibiting T-cell activation, proliferation, cytokine production and development of cytotoxicity. When expressed on the cell surface of tumor macrophages, plays an important role, together with regulatory T-cells (Treg), in the suppression of tumor-associated antigen-specific T-cell immunity. It also involved in promoting epithelial cell transformation.

### For Research Use Only