

Recombinant Human PDGF-AA Protein (His Tag)

Catalog Number: PKSH032905

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|----------------------|--|
| Species | Human |
| Source | E.coli-derived Human PDGF-AA protein Ser87-Thr 211, with an N-terminal His |
| Calculated MW | 15.9 kDa |
| Observed MW | 16 kDa |
| Accession | P04085 |
| Bio-activity | Not validated for activity |

Properties

| | |
|-----------------------|--|
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution of 4mM HCl. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Platelet-derived growth factor subunit A (PDGFA); belongs to the PDGF/VEGF growth factor family. PDGFA is a secreted protein; stored in platelet alpha-granules and released by platelets upon wounding. PDGFA is potent mitogens for a variety of cell types including smooth muscle cells; connective tissue cells; bone and cartilage cells; and some blood cells. It plays an essential role in the regulation of embryonic development; cell proliferation; cell migration; survival and chemotaxis. PDGFA is required for normal lung alveolar septum formation during embryogenesis; normal development of the gastrointestinal tract; normal development of Leydig cells and spermatogenesis; normal oligodendrocyte development and normal myelination in the spinal cord and cerebellum. It plays an important role in wound healing; Signaling is modulated by the formation of heterodimers with PDGFB.

For Research Use Only