A Reliable Research Partner in Life Science and Medicine

# Recombinant Human Interleukin-33/IL-33 Protein (His Tag)

Catalog Number: PKSH032642

Note: Centrifuge before opening to ensure complete recovery of vial contents.

)es						
 00	01	OH:	•	4-1	a	77
	w.		ш	ш	w	

Species Human

Source E.coli-derived Human Interleukin-33; IL-33 protein Serl 12-Thr 270, with an C-terminal

His

 Mol\_Mass
 18.9 kDa

 Accession
 O95760

**Bio-activity** Measure by its ability to binding with recombinant ST2L (IL-33 receptor). The ED<sub>50</sub>

for this effect is <54 ng/mL. Measure by its ability to induce proliferation in D10.G4.1 cells. The ED $_{50}$  for this effect is < 0.1 ng/mL. The specific activity of recombinant

human IL-33 is approximately >4 x10<sup>7</sup> IU/ mg.

## **Properties**

**Purity** > 98 % as determined by reducing SDS-PAGE.

**Endotoxin** < 0.1 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4.


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



> 98 % as determined by reducing SDS-PAGE.

## Background

Interleukin-33 (IL-33) belongs to the IL-1 superfamily. IL33 is highly expressed in endothelial venules found in tonsils; Peyer patches and mesenteric lymph nodes; but almost undetectable in placenta. IL33 induces the production of T helper-2 (Th2)-associated cytokines. IL-33 is a cytokine that mediates its biological effects by binding to and signals through IL1RL1/ST2 and IL-1 Receptor Accessory Protein (IL1RAP); activating intracellular molecules in the NF-κB and MAP kinase signaling pathways that drive production of type 2 cytokines from polarized Th2 cells.

## For Research Use Only

Fax: 1-832-243-6017