

# IL-17C Polyclonal Antibody(Capture/Detector)

catalog number: AN003970P

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description


<b>Reactivity</b>	Mouse;Human
<b>Immunogen</b>	Recombinant Mouse IL-17C Protein expressed by E.coli
<b>Host</b>	Rabbit
<b>Isotype</b>	Rabbit IgG
<b>Purification</b>	Antigen Affinity Purification
<b>Conjugation</b>	Unconjugated
<b>buffer</b>	Phosphate buffered solution, pH 7.2, containing 0.05% proclin 300.

## Applications

## Recommended Dilution

<b>ELISA Capture</b>	2-8 µg/mL
<b>ELISA Detector</b>	0.1-0.4 µg/mL

## Data



Sandwich ELISA-Recombinant Mouse IL-17C Protein standard curve. Background subtracted standard curve using Anti-IL-17C antibody(AN003970P)(Capture), Anti-IL-17C antibody(AN003970P)(Detector). The reference range value is 125-8000pg/mL for mouse.

## Preparation & Storage

<b>Storage</b>	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

## Background

The Interleukin 17 (IL-17) family proteins, comprising six members (IL-17, IL-17B through IL-17F), are secreted, structurally related proteins that share a conserved cysteine-knot fold near the C-terminus, but have considerable sequence divergence at the N-terminus. With the exception of IL-17B, which exists as a non-covalently linked dimer, all IL-17 family members are disulfide-linked dimers. IL-17 family proteins are pro-inflammatory cytokines that induce local cytokine production and are involved in the regulation of immune functions. Two receptors (IL-17 R, and IL-17B R), which are activated by IL-17 family members have been identified. In addition, at least three additional orphan type I transmembrane receptors with homology to IL-17 R, including IL-17 RL (IL-17 RC), IL-17 RD, and IL-17 RE, have also been reported. Mouse IL-17C cDNA encodes a 194 amino acid (aa) protein with a putative 14 aa signal peptide. Although there are no potential N-linked glycosylation sites, it is reportedly glycosylated. IL-17C shares from 15%-30% aa sequence identity with other IL-17 family members. Mouse and human IL-17C share 83% aa sequence identity. IL-17C has a very restricted expression pattern and was detected as a rare expressed sequence tag (EST) in an adult prostate and fetal kidney libraries. IL-17C has been shown to stimulate the release of TNF-alpha and IL-1 beta from the monocytic cell line THP-1, a property it shares with IL-17B. Human IL-17C is active on mouse cells.

## For Research Use Only