# **Elabscience**®

# Human IGF2R Antibody Pair Set

Catalog No.E-KAB-0252ApplicationsELISASynonymsCD222, CIMPR, M6P-R, MPR1, MPRI, cation-independent mannose-6-phosphate receptor

## Kit components & Storage

Title	Specifications	Storage
Human IGF2R Capture Antibody	1 vial, 100 µ g	Store at $-20^{\circ}$ C for one year.
		Avoid freeze / thaw cycles.
Human IGF2R Detection Antibody	1 vial, 50 μL	Store at $-20^{\circ}$ C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Product Information**

Items		Characteristic (E-KAB-0252)	
		Human ICE2D Conturn Antihody	Human IGF2R Detection Antibody
		Human IGF2R Capture Antibody	(Biotin)
Immunogen	Immunogen	Recombinant Human IGF2R protein	Recombinant Human IGF2R protein
Information	Swissprot	P11717	
Product details	Reactivity	Human	Human
	Host	Mouse	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%
		glycerol, pH 7.4	protective protein, 50% glycerol, pH
			7.4
	Purify	Protein A or G	Antigen Affinity
	Specificity	Detects Human IGF2R in ELISAs.	

For Research Use Only

# **Elabscience**®

# Applications

Human IGF2R Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4µg/mL	Human IGF2R Capture Antibody	
Capture			
ELISA	1:1000-1:10000	Human IGF2R Detection Antibody	Optical Density
Detection		(Biotin)	<sup>1</sup> 0.01 0.01 0.1 1 1 1 1 1 1 1 1 10 100 10

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

## Background

This gene encodes a receptor for both insulin-like growth factor 2 and mannose 6-phosphate, although the binding sites for either are located on different segments of the receptor. This receptor functions in the intracellular trafficking of lysosomal enzymes, the activation of transforming growth factor beta, and the degradation of insulin-like growth factor 2. While the related mouse gene shows exclusive expression from the maternal allele, imprinting of the human gene appears to be polymorphic, with only a minority of individuals showing expression from the maternal allele.

#### For Research Use Only