

## Recombinant Human Syntaxin-7/STX7 Protein (His Tag)

Catalog Number: PKSH033093

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human Syntaxin-7;STX7 protein Ser2-Leu238, with an N-terminal His
<b>Calculated MW</b>	28.0 kDa
<b>Observed MW</b>	31-36 kDa
<b>Accession</b>	O15400
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

Syntaxin 7 (STX7) is a member of a family of membrane-trafficking proteins named as soluble N-ethylmaleimide-sensitive factor attachment protein receptors (SNAREs), and mediates the endocytic trafficking from early endosomes to late endosomes and lysosomes. Syntaxin-7 has been shown to be present in both late endosomes and lysosomes, and to be required for both homotypic late endosome fusion and heterotypic fusion with lysosomes. STX7 has been shown to interact with STX8, VPS18, Vesicle-associated membrane protein 8 and VPS11.

### For Research Use Only