Recombinant Human Caspase-10/CASP10 Protein (His Tag)

Catalog Number: PKSH032176

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	E.coli-derived Human Caspase-10;CASP10 protein Val220-Ile480, with an C-terminal
	His
Calculated MW	30.1 kDa
Observed MW	33 kDa
Accession	Q92851-4
Bio-activity	Not validated for activity
Properties	
Purity	> 70 % as determined by reducing SDS-PAGE.
Concentration	Subject to label value.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at $<$ -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel
	packs. Upon receipt, store it immediately at $< -20^{\circ}$ C.
Formulation	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 8% Sucrose, 1mM DTT,
	0.05% Tween 80, pH8.5.
Data	
kDa	MK
120 90	
60	ence

> 70 % as determined	ed by reducing SDS-PAGE.

40 30

20

14

Background

Caspase-10 (CASP10) is a 521 amino acid protein member of the Cysteine-Aspartic Acid Protease (Caspase) family. CASP10 contains two DED (Death Effector) domains and is detectable in most tissues. CASP10 cleavage by Granzyme B and autocatalytic activity generate the two active subunits: Caspase-10 subunit p23/17, Caspase-10 subunit p12. Caspases are a family of cytosolic aspartate-specific cysteine proteases involved in the execution-phase of cell apoptosis, the initiation and execution. Human caspases can be subdivided into three functional groups: cytokine activation (caspase-1, -4, -5, and -13), apoptosis initiation (caspase-2, -8, -9, -and -10), and apoptosis execution (caspase-3, -6, and -7). CASP10 cleaves and activates caspases 3 and 7, but itself is processed by caspase 8. Defects in CASP10 are associated with apoptosis defects seen in type II autoimmune lymphoproliferative syndrome.

Web:www.elabscience.com