



A Reliable Research Partner in Life Science and Medicine

Human CCL22 Antibody Pair Set

Catalog No. E-KAB-0268 Applications ELISA

Synonyms CCL22, ABCD-1, DC/B-CK, SCYA22, STCP-1

Kit components & Storage

Title	Specifications	Storage
Human CCL22 Capture Antibody	1 vial, 100 μ g	Store at -20°C for one year.
		Avoid freeze / thaw cycles.
Human CCL22 Detection Antibody	1 vial, 50 μL	Store at -20°C for one year.
(Biotin)		Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Product Information

Items		Characteristic (E-KAB-0268)		
		Human CCL22 Capture Antibody	Human CCL22 Detection Antibody	
			(Biotin)	
Immunogen	Immunogen	Recombinant Human CCL22 protein	Recombinant Human CCL22 protein	
Information	Swissprot	O00626		
Product details	Reactivity	Human	Human	
	Host	Mouse	Chicken	
	Conjugation	Unconjugated	Biotin	
	Concentration	0.5mg/mL	/	
	Buffer	PBS with 0.04% Proclin 300, 50%	PBS with 0.04% Proclin 300, 1%	
		glycerol, pH 7.4	protective protein, 50% glycerol, pH	
			7.4	
	Purify	Protein A or G	Antigen Affinity	
	Specificity	Detects Human CCL22 in ELISAs.		

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com



A Reliable Research Partner in Life Science and Medicine

Applications

Human CCL22 Sandwich ELISA Assay:

	Recommended	Reagent	Images
	Concentration/Dilution		
ELISA	0.5-4μg/mL	Human CCL22 Capture Antibody	
Capture			Aist
ELISA Detection	1:1000-1:10000	Human CCL22 Detection Antibody (Biotin)	0.01 100 1000 10000 100000 Human MDC/CCL22 concentration(pg/mL)

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

The C-C motif chemokine ligand 22 (CCL22), also known as MDC, belongs to the group of chemokines, that are both constitutively expressed under homeostatic conditions and inducible upon inflammation. CCL22 is a secreted protein that exerts chemotactic activity for monocytes, dendritic cells, natural killer cells, and for chronically activated T lymphocytes. CCL22 is induced by LPS, IL-4, and IL-13 and in T cells by TCR stimulation. Accumulating studies indicate that CCL22 recruits regulatory T (T-reg) cells into tumor tissues and is expressed in many human tumors.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web: www.elabscience.com Email: techsupport@elabscience.com